


NONLINEAR OPTICS (PHYC/ECE 568)

Spring 2022 - Instructor: M. Sheik-Bahae

University of New Mexico

Homework #2, Due: Wed. Feb. 16

Problem 1. Two optical beams E_1 and E_2 with wavelengths of 1.0 and 0.6 μm respectively are incident on a nonlinear material.



(a) Assuming a $\chi^{(2)}$ nonlinearity, what new wavelengths can possibly be generated in this material?

The above nonlinear material is now replaced with a centro-symmetric material for the remaining part of this problem.

(b) What is the dominant nonlinear susceptibility?

(c) Assuming $\chi^{(3)}$ nonlinearity, what new wavelengths λ_j can possibly be generated in this material that simultaneously involve the interaction of both E_1 and E_2 beams? Write down the corresponding nonlinear polarization $P(\lambda_j)$ including their $\chi^{(3)}(\lambda_j; \lambda_k, \lambda_q, \lambda_p)$ terms (ignore Cartesian indices).

(d) If $|E_1| \gg |E_2|$, identify the most dominant terms in part (c).

(e) Write down the nonlinear polarization terms associated with self- and cross phase modulation of each beam (identify $\chi^{(3)}(\lambda_j; \lambda_k, \lambda_q, \lambda_p)$ terms)

(f) Under what condition the simultaneous presence of both beams leads to a nonlinear attenuation (absorption) of both beams? Describe this process, the required energy resonance (use diagrams), and the nature of the complex susceptibility $\chi^{(3)}(\lambda_j; \lambda_k, \lambda_q, \lambda_p)$ (with respect to part e).

(g) Under what condition the simultaneous presence of both beams leads to a nonlinear attenuation (absorption) of one beam (which?) and gain in the other (which?)? Describe this process, the required energy resonance (use diagrams), and the nature of the complex susceptibility $\chi^{(3)}(\lambda_j; \lambda_k, \lambda_q, \lambda_p)$ (with respect to part e and f).

Problem 2. Two-Photon Spectroscopy:

The 1S-2S transition in atomic Hydrogen ($E=10.206$ eV) is investigated using two-photon spectroscopy with two narrow-band CW laser sources. A pump laser with fixed wavelength $\lambda_1=200\text{nm}$ and a tunable laser ($\lambda=250\text{-}350$ nm) are used in a counter propagating arrangement as shown.

- (a) Qualitatively plot the transmission of the probe beam as a function of its tunable wavelength λ_2 .
- (b) Will the result in (a) be any different if they two beams were co-propagating? (Hint: think Doppler!)

